



# The results are in.

Patient self-testing and self-management through the use of a home INR monitor may be used to improve the time in therapeutic range (TTR) for select groups of patients. Increased TTR leads to improved clinical outcomes and reductions in thromboembolic and hemorrhagic events.<sup>1</sup>

## SAMPLE PATIENT RESULT

### ACS – PT/INR Self-Testing Service

Phone: (800) 769-8177  
Fax: (844) 700-3484

#### Patient Information


John Doe  
123 Main Street  
Anywhere, USA 55555

Last 4 SSN: 2356  
Home: (555) 555-5555

DOB: 11/13/1965  
Diagnosis: Atrial Fibrillation/I48.91  
Long-term use of anticoagulants/Zn9.01

INR Result: 4.1  
INR Range: 2-3  
Date Reported: 1/10/2016

**HIGH**



● Result ▼ Low Range ▲ High Range

Date	INR Range	INR
12/06/15	2 – 3	1.60
12/13/15	2 – 3	2.70
12/20/15	2 – 3	2.30
12/27/15	2 – 3	2.70
01/03/16	2 – 3	3.40
01/10/16	2 – 3	4.10

#### Clinician Information

Dr. Smith  
111 Main Street  
Suite 111  
Anytown, USA 55555

Contact: John  
Phone: (555) 555-5555  
Fax: (555) 555-5555

*The INR values in this report appear as reported by the patient to ACS.*

1. Pub 100-3 Medicare National Coverage Determinations, Chapter 1, Part 3, 190.11 – Home Prothrombin Time/International Normalized Ratio (PT/INR) Monitoring for Anticoagulation Management – Effective March 19, 2008 (Rev. 90, Issued: 07-25-08, Effective: 03-19-08, Implementation: 08-25-08)